

COHOMOLOGY, FUSION AND A P-NILPOTENCY CRITERION

JON GONZÁLEZ-SÁNCHEZ

ABSTRACT. Let G be a finite group, p a fix prime and P a Sylow p -subgroup of G . In this short note we prove that if p is odd, G is p -nilpotent if and only if P controls fusion of cyclic groups of order p . For the case $p = 2$, we show that G is p -nilpotent if and only if P controls fusion of cyclic groups of order 2 and 4.

1. INTRODUCTION

Throughout the text let p denote a fix prime. Let G be a finite group and P a Sylow p -subgroup of G . We denote by $H^\bullet(G, \mathbb{F}_p)$ the mod p cohomology algebra. It is well known that the restriction map in cohomology

$$(1) \quad H^\bullet(G, \mathbb{F}_p) \hookrightarrow H^\bullet(P, \mathbb{F}_p)$$

is injective (see [2, Proposition 4.2.2]). Suppose that G is p -nilpotent, i.e., P has a normal complement N in G . In this situation the composition

$$(2) \quad P \longrightarrow G \longrightarrow G/N,$$

is an isomorphism. Therefore the composition

$$(3) \quad H^\bullet(G/N, \mathbb{F}_p) \xrightarrow{\inf_{G/N}^G} H^\bullet(G, \mathbb{F}_p) \xrightarrow{\text{res}_P^G} H^\bullet(P, \mathbb{F}_p)$$

is also an isomorphism. This together with (1) implies that, if G is p -nilpotent, then the restriction map in cohomology $\text{res}_P^G : H^\bullet(G, \mathbb{F}_p) \longrightarrow H^\bullet(P, \mathbb{F}_p)$ is an isomorphism. The following result of M. Atiyah shows that the converse is also true.

Theorem 1 (Atiyah). *If $\text{res}_P^G : H^i(G, \mathbb{F}_p) \longrightarrow H^i(P, \mathbb{F}_p)$ are isomorphisms for all i big enough, then G is p -nilpotent. In particular G is p -nilpotent if and only if $\text{res}_P^G : H^\bullet(G, \mathbb{F}_p) \longrightarrow H^\bullet(P, \mathbb{F}_p)$ is an isomorphism.*

Proof. A proof of this can be found in the introduction of [8]. $\square$

Atiyah p -nilpotency criterion uses the cohomology in high dimension. Another cohomological criterion for p -nilpotency using cohomology in dimension 1 was provided by J. Tate ([10]).

Theorem 2 (Tate). *If $\text{res}_P^G : H^1(G, \mathbb{F}_p) \longrightarrow H^1(P, \mathbb{F}_p)$ is an isomorphism, then G is p -nilpotent.*

Proof. See [10]. $\square$

D. Quillen generalized Atiyah's p -nilpotency criterion for odd primes ([8]).

Theorem 3 (Quillen). *Let p be an odd prime. Then G is p -nilpotent if and only if $\text{res}_P^G : H^\bullet(G, \mathbb{F}_p) \longrightarrow H^\bullet(P, \mathbb{F}_p)$ is an F -isomorphism.*

Proof. See [8]. $\square$

Date: April 16, 2009.

2000 Mathematics Subject Classification. Primary 20D15, secondary 20J06.

Atiyah's p -nilpotency criterion can be reinterpreted in terms of p -fusion. We recall that a subgroup H of G *controls p -fusion in G* if

- (a) H contains a Sylow p -subgroup of G and
- (b) for any subgroup A of G and for any $g \in G$ such that $A, A^g \leq H$, there exists $x \in H$ such that for all $a \in A$, $a^g = a^x$.

By a result of G. Mislin [7], a subgroup H of G controls p -fusion in G if and only if $\text{res}_P^G : H^\bullet(G, \mathbb{F}_p) \longrightarrow H^\bullet(H, \mathbb{F}_p)$ is an isomorphism. Using Mislin's result Atiyah's p -nilpotency criterion follows from Frobenius p -nilpotency criterion.

Mislin's type of result can also be provided for the concept of F -isomorphism. In order to do this we introduce the following concept. Let $\mathcal{C}$ be a class of finite p -groups. We say that a subgroup H of G *controls fusion of $\mathcal{C}$ -groups in G* if

- (a) Any $\mathcal{C}$ -subgroup of G is conjugated to a subgroup of H and
- (b) for any $\mathcal{C}$ -subgroup A of G and for any $g \in G$ such that $A, A^g \leq H$, there exists $x \in H$ such that for all $a \in A$, $a^g = a^x$.

The condition (b) can be rewritten as

- (b') if A is a $\mathcal{C}$ -subgroup of H and $g \in G$ satisfies that $A^g \leq H$, then $g \in C_G(A).H$.

Theorem A below, which will be proved in Section 2, follows naturally from Quillen's work on cohomology (see [9] and [8]). Note that the "if" was proved in [4] and it is a direct consequence of Quillen's stratification ([9]). The converse follows from a careful reading of [8, Section 2].

Theorem A. *Let G be a finite group and H a subgroup of G . Then $\text{res}_H^G : H^\bullet(G, \mathbb{F}_p) \longrightarrow H^\bullet(H, \mathbb{F}_p)$ is an F -isomorphism if and only if H controls fusion of elementary abelian p -subgroups of G .*

In Section 3 we will prove the following p -nilpotency criterion that can be seen as a generalization of Quillen p -nilpotency criterion (Theorem 3 above) to the prime $p = 2$.

Theorem B. *Let G be a finite group and P a Sylow p -subgroup of G . Then the following two conditions are equivalent*

- (1) G is p -nilpotent.
- (2) P controls fusion of cyclic subgroups of order p in case p is odd, and cyclic subgroups of order 2 and 4 in case $p = 2$.

Note that Theorem A and Theorem B imply Quillen's p -nilpotency criterion. We will finish this short note by giving two applications of Theorem B. The first application will consist on reproving a result of H-W. Henn and S. Priddy that implies that "most" finite groups are p -nilpotent (see [5]). The second application is a generalization to the prime $p = 2$ of the following fact: if all elements of order p of a finite group G are in some upper center of G and p is an odd prime, then G is p -nilpotent (see [12] and [4]). For the prime $p = 2$ we will show that if all elements of order 2 and 4 are in some upper center of G , then G is 2-nilpotent.

We would like to end this introduction with an example of Quillen [8] where the necessity of considering cyclic groups of order 2 and 4 for the case $p = 2$ in Theorem B is illustrated.

Example 4. Consider $Q = \{1, -1, i, -i, j, -j, k, -k\}$ the quaternion group and α an automorphism of order 3 that permutes i, j and k . Let G be the semidirect product between Q and $\langle \alpha \rangle$ given by the action of α in Q . $A = \{1, -1\}$ is the only subgroup of exponent 2 in G . Clearly Q controls fusion of cyclic subgroup of order 2. However G is not 2-nilpotent.

2. COHOMOLOGY AND FUSION

The aim of this section is to sketch the proof of Theorem A. In subsections 2.1, 2.2 and 2.3 we will recall Quillen work in the mod p cohomology algebra of a finite group. This will be used in subsection 2.4 to prove Theorem A.

For a finite group G the *mod p cohomology algebra*

$$(4) \quad H^\bullet(G) = H^\bullet(G, \mathbb{F}_p)$$

is a finitely generated, connected, anti-commutative, $\mathbb{N}_0$ -graded $\mathbb{F}_p$ -algebra.

Let $\alpha_\bullet: A_\bullet \rightarrow B_\bullet$ be a homomorphism of finitely generated, connected, anti-commutative, $\mathbb{N}_0$ -graded $\mathbb{F}_p$ -algebras. Then $\alpha_\bullet$ is called an *F-isomorphism* if $\ker(\alpha_\bullet)$ is nilpotent, and for all $b \in B_n$ there exists $k \geq 0$ such that $b^{p^k} \in \text{im}(\alpha_\bullet)$.

2.1. Quillen's stratification. Let G be a finite group. Let $\mathcal{E}_G$ denote the category whose objects are the elementary abelian p -subgroups of G and whose morphisms are given by conjugation, i.e., for $E, E' \in \text{ob}(\mathcal{E}_G)$ one has

$$(5) \quad \text{mor}_G(E, E') = \{ i_g: E \rightarrow E' \mid g \in G, g E g^{-1} \leq E' \},$$

where $i_g(e) = g e g^{-1}$, $e \in E$. Then

$$(6) \quad H^\bullet(\mathcal{E}_G) = \varprojlim_{\mathcal{E}_G} H^\bullet(E)$$

is a finitely generated, connected, anti-commutative, $\mathbb{N}_0$ -graded $\mathbb{F}_p$ -algebra. Moreover, the restriction maps res_E^G yield a map

$$(7) \quad q_G = \prod_{E \in \text{ob}(\mathcal{E}_G)} \text{res}_E^G: H^\bullet(G) \longrightarrow H^\bullet(\mathcal{E}_G).$$

The following result is known as Quillen stratification.

Theorem 5 (Quillen). *Let G be a finite group. Then $q_G: H^\bullet(G) \rightarrow H^\bullet(\mathcal{E}_G)$ is an F-isomorphism.*

Proof. See [1, Cor. 5.6.4] or [9]. □

2.2. Cohomology of elementary abelian p -groups. One can easily deduce the cohomology of an elementary abelian p -group from the cohomology of the cyclic group of exponent p and the Kunneth formula.

Lemma 6. *Let A be an elementary abelian p -group. Then*

$$(8) \quad H^\bullet(A, \mathbb{F}_p) \cong \begin{cases} \Lambda(A^*) \otimes S(\beta(A^*)) & \text{if } p \text{ is odd} \\ S(A^*) & \text{if } p = 2, \end{cases}$$

where Λ denotes the exterior algebra functor, S the symmetric algebra functor, $A^* = \text{Hom}(A, \mathbb{F}_p) = H^1(A, \mathbb{F}_p)$ and β the Bockstein homomorphism from $H^1(A, \mathbb{F}_p)$ to $H^2(A, \mathbb{F}_p)$.

Proof. See [2, Chap. 3 Section 5]. □

From the previous lemma one can easily deduces that

$$(9) \quad H^\bullet(A, \mathbb{F}_p)/\sqrt{0} \cong S(A^*).$$

2.3. The spectrum of $H(G)$. Let G be a finite group. Following Quillen ([8]) we define

$$(10) \quad H(G) = \begin{cases} \oplus_{i \geq 0} H^{2i}(G, \mathbb{F}_p) & \text{if } p \text{ is odd} \\ \oplus_{i \geq 0} H^i(G, \mathbb{F}_p) & \text{if } p = 2. \end{cases}$$

$H(G)$ is a graded commutative ring. For an elementary abelian p -subgroup A of G , denote by $\mathfrak{g}_A$ the ideal of $H(G)$ consisting of elements u such that $u|_A$ is nilpotent. From (9), $\text{res}_A^G : H(G) \rightarrow H(A)$ induces a monomorphism

$$(11) \quad H(G)/\mathfrak{g}_A \hookrightarrow S(A^*).$$

In particular, the ideal $\mathfrak{g}_A$ is a prime ideal of $H(G)$. Furthermore,

Theorem 7 (Quillen). *Let $A, A' \subset G$ be elementary abelian subgroups of G . Then $\mathfrak{g}_A \subseteq \mathfrak{g}_{A'}$ if and only if A' is conjugated to a subgroup of A . In particular $\mathfrak{g}_A = \mathfrak{g}_{A'}$ if and only if A and A' are conjugated in G .*

Proof. See [8, Theorem 2.7]. $\square$

Let us consider the extension of quotient fields associated to the monomorphism in (11),

$$(12) \quad k(\mathfrak{g}_A) \hookrightarrow k(A).$$

We have that

Theorem 8 (Quillen). *The extension $k(A)/k(\mathfrak{g}_A)$ is a normal extension and*

$$(13) \quad \text{Aut}(k(A)/k(\mathfrak{g}_A)) \cong N_G(A)/C_G(A).$$

Proof. See [8, Theorem 2.10]. $\square$

2.4. F-isomorphisms and fusion. The following lemma is a standard result in commutative algebra.

Lemma 9. *Let A and B be commutative $\mathbb{F}_p$ -algebras and $f : A \rightarrow B$ an F -isomorphism. Then $f^* : \text{Spec}(B) \rightarrow \text{Spec}(A)$ is a homeomorphism.*

Proof. Since the kernel of f is nilpotent, then for any radical ideal $\mathfrak{a}$ of A one has that $f^{-1}(\sqrt{f(\mathfrak{a})}) = \mathfrak{a}$. Since for any $x \in B$ there exists $y \in A$ and $n \geq 0$ such that $f(y) = x^{p^n}$, then for any radical ideal $\mathfrak{b}$ of B one has that $\sqrt{f(f^{-1}(\mathfrak{b}))} = \mathfrak{b}$. Therefore

$$(14) \quad \mathfrak{a} \longrightarrow \sqrt{f(\mathfrak{a})}$$

$$(15) \quad \mathfrak{b} \longrightarrow f^{-1}(\mathfrak{b})$$

is a bijection between the radical ideals of A and the radical ideals of B . In particular f^* is an isomorphism of varieties. $\square$

We are now ready to prove Theorem A.

Proof of Theorem A. Suppose first that H controls fusion of elementary abelian p -subgroups of G . Then the embedding functor

$$(16) \quad j_{H,G} : \mathcal{E}_H \longrightarrow \mathcal{E}_G$$

is an equivalence of categories. Therefore

$$(17) \quad H^\bullet(j_{H,G}) : H^\bullet(\mathcal{E}_G) \longrightarrow H^\bullet(\mathcal{E}_H)$$

is an isomorphism. Consider the commutative diagram

$$(18) \quad \begin{array}{ccc} H^\bullet(G) & \xrightarrow{q_G} & H^\bullet(C_G) \\ \text{res}_H^G \downarrow & & \downarrow H^\bullet(j_{H,G}) \\ H^\bullet(H) & \xrightarrow{q_H} & H^\bullet(C_H). \end{array}$$

By Theorem 5 and equation (17) it follows that res_H^G is an F -isomorphism.

Suppose now that $\text{res}_H^G: H^\bullet(G, \mathbb{F}_p) \rightarrow H^\bullet(H, \mathbb{F}_p)$ is an F -isomorphism. Then res_H^G induces an F -isomorphism $f: H(G) \rightarrow H(H)$. Consider A and A' two elementary abelian p -subgroups of H .

Subclaim 1: If A and A' are conjugated in G , then they are conjugated in H .

Subproof. By Lemma 9, $f^*: \text{Spec}(H(H)) \rightarrow \text{Spec}(H(G))$ provides a bijection between the prime ideals of $H(H)$ and the prime ideals of $H(G)$. Furthermore, if A is an elementary abelian p -subgroup of H , then $\mathfrak{g}_A = f^*(\mathfrak{h}_A)$. By Theorem 7, if A and A' are conjugated in G , then $\mathfrak{g}_A = \mathfrak{g}_{A'}$. In particular $f^*(\mathfrak{h}_A) = \mathfrak{g}_A = \mathfrak{g}_{A'} = f^*(\mathfrak{h}_{A'})$. Therefore $\mathfrak{h}_A = \mathfrak{h}_{A'}$. Hence, by Theorem 7, A and A' are conjugated in H . $\square$

Subclaim 2: $N_G(A) = C_G(A)N_H(A)$.

Subproof. Since $k(\mathfrak{h}_A)$ is a purely inseparable extension of $k(\mathfrak{g}_A)$, then

$$(19) \quad \text{Aut}(k(A)/k(\mathfrak{h}_A)) \cong \text{Aut}(k(A)/k(\mathfrak{g}_A)).$$

Therefore, by Theorem 8, $N_H(A)/C_H(A) \cong N_G(A)/C_G(A)$. $\square$

Subclaim 3: H controls fusion of elementary abelian p -subgroups of G .

Subproof: Let A be an elementary abelian p -subgroup of H and $g \in G$ such that $A^g \leq H$. Then, by Subclaim 1 there exists $h \in H$ such that $A^g = A^h$. In particular, by Subclaim 2, $gh^{-1} \in N_G(A) = C_G(A).N_H(A)$. Therefore $g \in C_G(A).H$ $\square$

3. A p -NILPOTENCY CRITERION

In this section we will prove our main result Theorem B. To ease the notation we denote by $\mathcal{C}_p$ the class of cyclic groups of order p in case p is odd and cyclic groups of order 2 and 4 in case $p = 2$. Put $\mathbf{p} = p$ if p is odd and $\mathbf{p} = 4$ in case $p = 2$.

Theorem 10. *Let G be a finite group and P a Sylow p -subgroup of G . Then the following two conditions are equivalent*

- (1) G is p -nilpotent.
- (2) P controls fusion of $\mathcal{C}_p$ -groups.

Proof. It is clear that if G is p -nilpotent, then P controls fusion of $\mathcal{C}_p$ -groups.

Let us show the converse. Using Frobenius p -nilpotency criterion it is enough to prove that for any subgroup B of P and for any p' -element $g \in N_G(B)$, then g centralizes B . The subgroup B is contained in $Z_l(P)$ for some $l \geq 1$ where $Z_l(P)$ denotes the l -upper center of P . We will show by induction on l that $g \in C_G(B)$. Suppose first that $B \leq Z(P)$ and consider $a \in B$ such that $a^{\mathbf{p}} = 1$. Since P controls fusion of $\mathcal{C}_p$ -groups, there exists $x \in P$ such that $a^g = a^x$ and since $a \in Z(P)$, then $a^x = a$. Hence we have that g centralizes all elements of order p (2 and 4 in case $p = 2$) in B . Thus, by [6, Chap. V Lemma 5.12], g centralizes B .

For the general case, consider $B \leq Z_l(P)$ and suppose the assumption to be true for any subgroup contained in $Z_{l-1}(P)$.

Subclaim 1: For $a \in B$ such that $a^{\mathbf{p}} = 1$, we have that $[a, g, g] = 1$.

Subproof. We have that g normalizes the subgroups $K = \langle a \in B \mid a^{\mathbf{p}} = 1 \rangle$ and $[K, g]$. We also have that

$$(20) \quad [K, g] = \langle [a, g]^b \mid a, b \in K \text{ and } a^{\mathbf{p}} = 1 \rangle.$$

Take $a \in B$ such that $a^{\mathbf{p}} = 1$. Since P controls fusion of $\mathcal{C}_{\mathbf{p}}$ -groups, there exists $x \in P$ such that $a^g = a^x$. In particular $[a, g] = [a, x] \in Z_{l-1}(P)$. Therefore, by (20), $[K, g] = Z_{l-1}(P)$. Since g normalizes $[K, g]$ and by induction hypothesis we have that $[K, g, g] = 1$. $\square$

Subclaim 2: $g \in C_G(B)$.

Subproof. Take $a \in B$ such that $a^{\mathbf{p}} = 1$ and put p^e the exponent of B . Consider the subgroup $H = \langle g, [a, g] \rangle$. By the Subclaim 1, $\gamma_2(H) = 1$. Then, by [6, Chap. III, Theorem 9.4], we have that $[a, g^{p^e}] = [a, g]^{p^e} = 1$. Since g is a p' -element of G , g centralizes all elements of order p (2 and 4 in case $p = 2$) in B . Thus, by [6, Chap. V Lemma 5.12], g centralizes B .

This ends the proof. $\square$

As a consequence to this we have the following corollary.

Corollary 11. *Let G a finite group and P a Sylow p -subgroup of G such that*

1. $N_G(P)$ controls fusion of $\mathcal{C}_{\mathbf{p}}$ -groups and
2. $N_G(P) = C_G(P).P$.

Then G is p -nilpotent.

Proof. Let A be a $\mathcal{C}_{\mathbf{p}}$ -group and $g \in G$ such that $A, A^g \leq P$. Since $N_G(P)$ controls fusion $\mathcal{C}_{\mathbf{p}}$ -groups, one has that $g \in C_G(A).N_G(P) = C_G(A).P$. Then P controls fusion of $\mathcal{C}_{\mathbf{p}}$ -groups and, by Theorem 10, G is p -nilpotent. $\square$

4. SOME APPLICATIONS

We now present the first application of Theorem 10. In [5] H-W. Henn and S. Priddy proved that if a group G has a Sylow p -subgroup P such that

- i) if p is odd, the elements of order p of P are in the center of P and, if $p = 2$, the elements of order 2 and 4 are in the center of P ,
- ii) $\text{Aut}(P)$ is a p -group,

then G is p -nilpotent. This implies that "most" finite groups are p -nilpotent (see [5]). The proof of Henn and Priddy is essentially topological. In [11] J. Thevenaz gave a group theoretical proof of this result using Alperin's Fusion Theorem. In fact Thevenaz proved that if G satisfies condition i), then $N_G(P)$ controls p -fusion in G . This, together with condition ii) above implies that P controls p -fusion in G and therefore G is p -nilpotent. We now give a weaker version of Thevenaz result which also implies that a group satisfying i) and ii) is p -nilpotent.

Proposition 12. *Let G be a finite group and P a Sylow p -subgroup of G . Suppose that the elements of order dividing p in P (or 4 in case $p = 2$) are in the center of P . Then $N_G(P)$ controls fusion of $\mathcal{C}_{\mathbf{p}}$ -groups.*

Proof. Let A be a $\mathcal{C}_{\mathbf{p}}$ -group and $g \in G$ such that $A, A^g \leq N_G(P)$. In particular $A^g \leq P$. Equivalently $A \leq P^{g^{-1}}$. Hence, since the elements of P of order p (or 4 in case $p = 2$) are in the center of P , we have that $P, P^{g^{-1}} \leq C_G(A)$. But, since P and $P^{g^{-1}}$ are Sylow p -subgroups of $C_G(A)$, there exists $c \in C_G(A)$ such that $P = P^{g^{-1}c}$. Thus $g^{-1}c \in N_G(P)$ and $g \in N_G(P).C_G(A)$. $\square$

Corollary 13. *Let G a finite group and P a Sylow p -subgroup of G such that*

1. all elements of order dividing p in P (or 4 in case $p = 2$) are in the center of P and
2. $N_G(P) = P.C_G(P)$.

Then G is p -nilpotent.

Proof. It follows from Proposition 12 and Corollary 11. $\square$

The second application of Theorem 10 is a generalization to $p = 2$ of the fact that if the elements of order p of a finite group G are in some upper center of G , then G is p -nilpotent (see [12] and [4]).

Corollary 14. *Let G a finite group such that $K = \langle x \in G \mid x^{\mathbf{p}} = 1 \rangle \leq Z_n(G)$ for some $n \geq 1$ (here $\mathbf{p}$ means p in case p is odd and 4 in case $p = 2$). Then G is p -nilpotent.*

Proof. The subgroup K is nilpotent of class at most n , and therefore a finite p -group. Let p^e be the exponent of K . Then, by Hall-Petrescu collection formula (see [3, Theorem 2.1]), for any $y \in K$ and $x \in G$

$$(21) \quad [y, x^{p^{e+n}}] \in \prod_{0 \leq i \leq e+n} [K, \overbrace{G, \dots, G}^{p^i}]^{p^{e+n-i}} = 1.$$

Therefore one has that $G^{p^{e+n}} \leq C_G(K)$. Moreover, for any Sylow p -subgroup P of G one has $G = P.G^{p^{e+n}} = P.C_G(K)$. In particular P controls fusion of $\mathcal{C}_{\mathbf{p}}$ -groups. Hence, by Theorem 10, G is p -nilpotent. $\square$

REFERENCES

- [1] D. J. Benson, *Representations and cohomology. II*, Cambridge Studies in Advanced Mathematics, vol. 31, Cambridge University Press, Cambridge, 1991, Cohomology of groups and modules.
- [2] L. Evens, *The cohomology of groups*, Oxford Mathematical Monographs, The Clarendon Press Oxford University Press, New York, 1991, Oxford Science Publications.
- [3] G. Fernández-Alcober, J. González-Sánchez, and A. Jaikin-Zapirain, *Omega subgroups of pro- p groups*, Israel J. Math. **166** (2008), 393–412.
- [4] J. González-Sánchez and T. Weigel, *p -central groups of height k* , preprint, 2008.
- [5] H.-W. Henn and S. Priddy, *p -nilpotence, classifying space indecomposability, and other properties of almost all finite groups*, Comment. Math. Helv. **69** (1994), no. 3, 335–350.
- [6] B. Huppert, *Endliche Gruppen. I*, Die Grundlehren der Mathematischen Wissenschaften, Band 134, Springer-Verlag, Berlin, 1967.
- [7] G. Mislin, *On group homomorphisms inducing mod- p cohomology isomorphisms*, Comment. Math. Helv. **65** (1990), no. 3, 454–461.
- [8] D. Quillen, *A cohomological criterion for p -nilpotence*, J. Pure Appl. Algebra **1** (1971), no. 4, 361–372.
- [9] ———, *The spectrum of an equivariant cohomology ring. I, II*, Ann. of Math. (2) **94** (1971), 549–572; ibid. (2) **94** (1971), 573–602.
- [10] J. Tate, *Nilpotent quotient groups*, Topology **3** (1964), no. suppl. 1, 109–111.
- [11] J. Thevenaz, *Most finite groups are p -nilpotent.*, Expo. Math. **11** (1993), no. 4, 359–363.
- [12] Th. S. Weigel, *p -central groups and Poincaré duality*, Trans. Amer. Math. Soc. **352** (2000), no. 9, 4143–4154.

JON GONZÁLEZ-SÁNCHEZ, UNIVERSIDAD DE CANTABRIA, DEPARTAMENTO DE MATEMÁTICAS, ESTADÍSTICA Y COMPUTACIÓN, FACULTAD DE CIENCIAS, AVDA. DE LOS CASTROS, s/n, E-39071 SANTANDER, SPAIN

E-mail address: jon.gonzalez@unican.es